

Hypoglycemia Emergency Care Plan (For Low Blood Glucose)

Student's Name: _____

Grade/Teacher: _____

Date of Plan: _____

Emergency contact information

Parent 1/Guardian: _____

Email Address: _____ Home Phone: _____

Work Phone: _____ Mobile: _____

Parent 2/Guardian: _____

Email Address: _____ Home Phone: _____

Work Phone: _____ Mobile: _____

Health Care Provider: _____

Phone Number: _____

School Nurse: _____

Contact Number(s): _____

Trained Diabetes Personnel: _____

Contact Number(s): _____

The student should never be left alone, or sent anywhere alone or with another student, when experiencing hypoglycemia.

Causes of Hypoglycemia	Onset of Hypoglycemia
<ul style="list-style-type: none"> • Too much insulin • Missing or delaying meals or snacks • Not eating enough food (carbohydrates) • Getting extra, intense, or unplanned physical activity • Being ill, particularly with gastrointestinal illness 	<ul style="list-style-type: none"> • Sudden—symptoms may progress rapidly

Hypoglycemia Symptoms		
Circle student's usual symptoms.		
Mild to Moderate		Severe
<ul style="list-style-type: none"> • Shaky or jittery • Sweaty • Hungry • Pale • Headache • Blurry vision • Sleepy • Dizzy • Lightheaded • Confused • Disoriented 	<ul style="list-style-type: none"> • Uncoordinated • Irritable or nervous • Argumentative • Combative • Changed personality • Changed behavior • Inability to concentrate • Weak • Lethargic • Other: _____ 	<ul style="list-style-type: none"> • Inability to eat or drink • Unconscious • Unresponsive • Seizure activity or convulsions (jerking movements)

Actions for Treating Hypoglycemia	
<p>Notify school nurse or trained diabetes personnel as soon as you observe symptoms. If possible, check blood glucose (sugar) at side of finger. Treat for hypoglycemia if blood glucose level is less than _____ mg/dL.</p> <p>WHEN IN DOUBT, ALWAYS TREAT FOR HYPOGLYCEMIA AS SPECIFIED BELOW.</p>	
Treatment for Mild to Moderate Hypoglycemia	Treatment for Severe Hypoglycemia
<input type="checkbox"/> Provide quick-acting glucose (sugar) product equal to _____ grams of carbohydrates. Examples of 15 grams of carbohydrates are listed below: <ul style="list-style-type: none"> • 4 glucose tablets • 1 tube of glucose gel • 4 ounces of fruit juice (not low-calorie or reduced-sugar) • 4–6 ounces (1/2 can) of soda (not low-calorie or reduced-sugar) <input type="checkbox"/> Wait 15 minutes. <input type="checkbox"/> Recheck blood glucose level. <input type="checkbox"/> Repeat quick-acting glucose product if blood glucose level is less than _____mg/dL. <input type="checkbox"/> Contact the student's parents/guardians. <input type="checkbox"/> Once the student's blood glucose returns to normal, check the blood glucose level 1 hour later. Provide an additional source of carbohydrate (e.g., whole grain crackers, graham crackers, granola bar, yogurt, or fruit) if a meal or snack is not planned.	<input type="checkbox"/> Position the student on his or her side. <input type="checkbox"/> Do not attempt to give anything by mouth. <input type="checkbox"/> Administer glucagon: _____ mg at _____ site. <input type="checkbox"/> While treating, have another person call 911 (Emergency Medical Services). <input type="checkbox"/> Contact student's parents/guardians. <input type="checkbox"/> Stay with student until Emergency Medical Services arrive. <input type="checkbox"/> Notify student's health care provider.